



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2018

---

**Complete and assembled genome sequence of *Salmonella enterica* subsp. *enterica* serotype Senftenberg N17-509, a strain lacking *Salmonella* pathogen island SPI-1**

Stevens, Marc J A ; Zurfluh, Katrin ; Althaus, D ; Corti, S ; Lehner, Angelika ; Stephan, Roger

**Abstract:** The genome of *Salmonella enterica* subsp. *enterica* serotype Senftenberg N17-509, a strain isolated from desiccated coconut, was sequenced using single-molecule real-time sequencing. It consists of a 5.1-Mbp chromosome and a 29-kb linear plasmid.

DOI: <https://doi.org/10.1128/genomeA.00156-18>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-167635>

Journal Article

Accepted Version

Originally published at:

Stevens, Marc J A; Zurfluh, Katrin; Althaus, D; Corti, S; Lehner, Angelika; Stephan, Roger (2018). Complete and assembled genome sequence of *Salmonella enterica* subsp. *enterica* serotype Senftenberg N17-509, a strain lacking *Salmonella* pathogen island SPI-1. *Genome Announcements*, 6(12):e00156-18.

DOI: <https://doi.org/10.1128/genomeA.00156-18>

**Complete and assembled genome sequence of *Salmonella enterica* subsp. *enterica***  
**serotype Senftenberg N17-509, a strain lacking *Salmonella* pathogen island SPI-1**

Marc J. A. Stevens, Katrin Zurfluh, Denise Althaus, Sabrina Corti, Angelika Lehner,  
Roger Stephan\*

Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich,  
Switzerland.

\*Corresponding author: roger.stephan@uhz.ch

**Abstract.**

The genome of *Salmonella enterica* subsp. *enterica* serotype Senftenberg N17-509, a strain isolated from desiccated coconut, was sequenced using single molecule real time sequencing. It consists of a 5.1-Mbp chromosome, and a 29-kb linear plasmid.

The *Salmonella enterica* subsp. *enterica* strain N17-509 was isolated from desiccated coconut and provided different identification and typing challenges. The biochemical tests performed to confirm the *Salmonella* genus were inconclusive. A commercial real-time PCR identification system (Assurance GDS® *Salmonella* kit (BioControl Systems, Bellevue, WA, USA)) based on the *invA* gene yielded a negative result. A further rRNA based fast method (VIT *Salmonella* kit (vermicon, München, Germany) as well as the

MALDI TOF based identification was positive for *Salmonella*. According to the KAUFFMANN-WHITE Scheme (Institut Pasteur, Paris), the strain was serotyped as *Salmonella enterica* subsp. *enterica* serovar 1,3,19:nt.

Genomic DNA was extracted using Wizard® Genomic DNA Purification Kit according to the manufacturer's protocol (Promega AG, Dübendorf, Switzerland). The genome was sequenced using a two SMRT cells on a PacBio RS II (Pacific Biosciences, Menlo Park, CA, USA) and was performed at the Functional Genomics Center Zurich (Zurich, Switzerland). The raw reads were filtered using the RS\_Filter\_Only protocol in the SMRT-portal (Pacific Biosciences) with 0.75 as polymerase read quality cut off and a minimal length of 500 bp. A total of 91,709 reads with an average length of 11,001 bp were selected, corresponding to 1,008,898,178 sequenced base pairs and a 197-fold coverage. The reads were assembled using the Canu assembler 1.6 (1) with the option "pacbio-raw" and an estimated genome size of 5.1 Mbp. The Canu output consisted of 3 contigs, which were further polished in CLC workbench 7 (CLC, Aarhus, Denmark). Two contigs were combined and formed the chromosomal sequence. The third contig contained a plasmid-like sequence. The start of the chromosome was set at the origin of replication, determined using OriCfinder (2).

A blastN with the complete chromosome against all *Enterobacteriales* genomes in the NCBI nucleotide database revealed highest identity of 99% and a coverage of 91% with the genome of *Salmonella enterica* subsp. *enterica* serotype Senftenberg 775W (=ATCC 43845). Hence, strain N17-509 was classified as *Salmonella enterica* subsp. *enterica* serotype Senftenberg. The genome of N17-509 consists of a 5,121,989-bp chromosome and a 29,600-bp extra-chromosomal element designated pN17-509. The GC-contents of

the chromosome and pN17-509 are 52.2% and 36,5%, respectively. The genome was annotated by the NCBI Prokaryotic Genomes Automatic Annotation Pipeline and contains 5361 genes of which 5241 encoding proteins. Further, 84 tRNA genes and 7 rRNA operons are present. Remarkably, the *Salmonella* pathogen island SPI-1, harbouring e.g. the *invA* gene, is absent in this strain. The plasmid pN17-509 has 97% identity at 87% coverage with the linear plasmid pBSSB1 from *Salmonella enterica* subsp. *enterica* serovar Typhi that mediates flagellar variation (3).

**Nucleotide sequence accession number.** Sequence and annotation data of the complete genome of *Salmonella enterica* subsp. *enterica* serotype Senftenberg N17-509 are deposited in the GenBank database with the accession number CP026379 for the chromosome and CP026380 for the plasmid.

## REFERENCES

- 1) Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* **27**:722-736.
- 2) Gao F, Zhang CT. 2008. Ori-Finder: a web-based system for finding oriCs in unannotated bacterial genomes. *BMC Bioinformatics* **9**:79.

69        3) **Baker S, Hardy J, Sanderson KE, Quail M, Goodhead I, Kingsley RA,**  
70            **Parkhill J, Stocker B, Dougan G.** 2007. A novel linear plasmid mediates  
71            flagellar variation in *Salmonella* Typhi. PLoS Pathog **11**;3(5):e59.